

Routine Immunization Screening Form: Adult

NOTE: If cholera or smallpox vaccines are being considered, please complete their respective immunization screening forms.

AUTHORITY: 10 U.S.C. 1071-1085, Medical and Dental Care; Army Regulation 40-562, Immunizations and Chemoprophylaxis for the Prevention of Infectious Disease; DoDM 6025.18, Implementation of the Health Insurance Portability and Accountability Act (HIPAA) Privacy Rule in DoD Health Care Programs.

PURPOSE: To determine whether you can safely receive a routine immunization.

ROUTINE USES: Use and disclosure of your records outside of DoD may occur in accordance with the Privacy Act of 1974, as amended (5 U.S.C. 552a(b)). Collected information may be shared with entities including the Departments of Health and Human Services, Veterans Affairs, and other Federal, State, local, or foreign government agencies, or authorized private business entities.

To appropriate agencies, entities, and persons when (1) the DoD suspects or has confirmed that there has been a breach of the system of records; (2) the DoD has determined that as a result of the suspected or confirmed breach there is a risk of harm to individuals, the DoD (including its information systems, programs, and operations), the Federal Government, or national security; and (3) the disclosure made to such agencies, entities, and persons is reasonably necessary to assist in connection with the DoD's efforts to respond to the suspected or confirmed breach or to prevent, minimize, or remedy such harm.

APPLICABLE SORN: EDHA 07, Military Health Information System (November 18, 2013, 78 FR 69076) <https://dpcl.d.defense.gov/Privacy/SORNsIndex/DOD-wide-SORN-Article-View/Article/570672/edha-07/>

DISCLOSURE: Voluntary. If you choose not to provide the requested information, no penalty may be imposed; however, failure to provide the information may result in delays in assessing contraindications for receiving vaccinations.

Patient name: _____ DOB (YYYYMMDD): _____

Screening Checklist for Contraindications to Vaccines for Adults

For patients: The following questions will help us determine which vaccines you may be given today. If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

		Yes	No	Don't Know
1.	Are you sick today?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2.	Have you ever had a serious reaction after receiving a vaccination?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3.	Do you have allergies to medication, food, a vaccine component, or latex?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4.	Have you had a seizure or a brain or other nervous system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5.	Have you had a health problem involving heart, lung (e.g., asthma), kidney, or metabolic disease (e.g., diabetes), anemia, or other blood disorder?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.	Do you have cancer, leukemia, HIV/AIDS, or any other immune system problem?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7.	In the past 3 months, have you taken medications that weaken your immune system, such as prednisone or other steroids; anticancer drugs; biologic drugs for autoimmune diseases such as rheumatoid arthritis, Crohn's disease, or psoriasis; or had radiation treatments?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8.	In the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9.	Have you had (or are you a candidate for) your spleen removed, or do you have sickle cell anemia?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10.	Have you ever passed out (had vasovagal syncope) during or after a previous immunization or blood draw?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11.	Have you received any vaccinations in the past 4 weeks?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
12.	Are you pregnant or is there a chance you could become pregnant during the next month?	Not Applicable <input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Please list any medications you are currently taking:

Form completed by: _____ Date (YYYYMMDD): _____

Form reviewed by: _____ Date (YYYYMMDD): _____

Did you bring your immunization record/card with you? Yes No

It is important for you to have a personal record of your vaccinations. If you don't have a personal record, ask your healthcare provider to give you one. Keep this record in a safe place and bring it with you every time you seek medical care. Make sure your healthcare provider records all your vaccinations on it. For questions or concerns regarding immunizations, providers, nurses and patients may call the DHA Immunization Healthcare Support Center 24/7 at 1-877-438-8222, Option 1.

Information for Healthcare Professionals about the Screening Checklist for Contraindications (Adult)

Each screening question is explained in more detail below. For more information, please consult the sources referenced at the end.

1. Are you sick today? [all vaccines]

There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events.¹ However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. Mild illnesses (such as otitis media, upper respiratory infections, and diarrhea) are NOT contraindications to vaccination. Do not withhold vaccination if a person is taking antibiotics.

2. Have you ever had a serious reaction after receiving a vaccination? [all vaccines]

History of anaphylactic reaction (see question 3) to a previous dose of vaccine or vaccine component is a contraindication for subsequent doses.⁴ History of encephalopathy within 7 days following DTP/DTaP is a contraindication for further doses of pertussis-containing vaccine. There are other adverse events that may occur following vaccination that constitute contraindications or precautions to future doses. Under normal circumstances, vaccines are deferred when a precaution is present. However, situations may arise when the benefit outweighs the risk (e.g., during a community pertussis outbreak).

3. Do you have allergies to medications, food, a vaccine component, or latex? [all vaccines]

An anaphylactic reaction to latex is a contraindication to vaccines that contain latex as a component or as part of the packaging (e.g., vial stoppers, prefilled syringe plungers or caps). If a person has anaphylaxis after eating gelatin, do not administer vaccines containing gelatin. For patients with known Alpha-gal syndrome (red meat allergy), caution should be exercised with gelatin-containing vaccines (i.e. MMR, VAR, YF-Vax), as some of these patients have demonstrated anaphylaxis with these vaccines. A local reaction to a prior vaccine dose or vaccine component, including latex, is not a contraindication to a subsequent dose or vaccine containing that component.^{2,3} People with egg allergy of any severity can receive any recommended influenza vaccine (i.e., any IIV or RIV) that is otherwise appropriate for the patient's age. Vaccine administration should be supervised by a healthcare provider who is able to recognize and manage severe allergic conditions.⁴

4. Have you had a seizure, or had brain or other nervous system problems?

[IIV, LAIV, Td, Tdap]

Tdap is contraindicated in patients who have a history of encephalopathy within 7 days following DTP/DTaP given as a child. An unstable, progressive neurologic condition is a precaution to the use of Tdap. For patients with stable neurologic disorders (including seizures) unrelated to vaccination, or for patients with a family history of seizures, vaccinate as usual. A history of Guillain-Barre syndrome (GBS) is a precaution for the following: 1) Td/Tdap: if GBS occurred within 6 weeks of a tetanus-containing vaccine and the decision is made to continue vaccination, if no history of prior Tdap, give Tdap instead of Td; 2) Influenza vaccine (IIV or LAIV): if GBS occurred within 6 weeks of a prior influenza vaccination, vaccinate with IIV if at high risk for severe influenza complications.

5. Have you had a health problem involving heart, lung (e.g., asthma), kidney, or metabolic disease (e.g., diabetes), anemia, or other blood disorder? [MMR, LAIV]

A history of thrombocytopenia or thrombocytopenic purpura is a precaution to MMR vaccine. The safety of LAIV in patients with these conditions has not been established. These conditions, including asthma in adults, should be considered precautions for LAIV use.

6. Do you have cancer, leukemia, HIV/AIDS, or any other immune system problem?

[Adenovirus, CVD 103-HgR, LAIV, MMR, Ty21a, VAR, YF vaccine, CHIKV]

Live virus vaccines are usually contraindicated in immunocompromised patients; however, there are exceptions. MMR is recommended and varicella should be considered for adults with CD4+ T-lymphocyte counts of greater than or equal to 200cell/ μ L. Immunosuppressed patients should not receive LAIV. For details, consult current ACIP recommendations.^{1,6,7,8}

7. In the past 3 months, have you taken medications that weaken your immune system, such as prednisone or other steroids; anticancer drugs; biologic drugs for autoimmune diseases such as rheumatoid arthritis, Crohn's disease, or psoriasis; or had radiation treatments? [Adenovirus, CVD 103-HgR, MMR, Ty21a, VAR, YF vaccine, CHIKV]

Live virus vaccines should be postponed until after chemotherapy or long-term, high-dose

steroid therapy has ended. For details and length of time to postpone, consult the current ACIP statement.¹ Some immune mediator and immune modulator drugs (especially the antitumor necrosis factor agents adalimumab, infliximab, and etanercept) may be immunosuppressive. The use of live vaccines should be avoided in persons taking these drugs.¹ Specific vaccination schedules for stem cell transplant (bone marrow transplant) patients can be found on the NIH website.⁸ LAIV, when recommended, can be given only to healthy, non-pregnant women ages 2 through 49 years.

8. In the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug?

[MMR, VAR]

Certain live virus vaccines may need to be deferred, depending on several variables. Consult the most current ACIP recommendations for information on intervals between receipt of antiviral drugs, immune globulin or blood products, and live virus vaccines.^{1,7}

9. Have you had (or are you a candidate for) your spleen removed, or do you have sickle cell anemia? [Hib, LAIV, PCV13, PPSV23, MenACWY, MenB]

Patients with anatomic or functional asplenia (i.e. sickle-cell disease) are at an increased risk of certain vaccine preventable diseases to include Haemophilus influenzae type b, meningococcal, and pneumococcal disease. LAIV is not recommended for people with anatomic or functional asplenia. Hib, PCV13, MCV4, and MenB vaccine should be given 14 days before splenectomy, if possible. Doses given during the 14 days prior to surgery can be counted as valid. Doses that cannot be given prior to surgery should be given as soon as the patient's condition has stabilized after surgery. For patients 2 years of age and up: the first dose of PPSV23 should be administered 8 weeks after the last dose of PCV13. A second dose of PPSV23 should be administered 5 years after the first dose. A third, final dose of PPSV23 should be administered after age 65 years, if both previous doses were before the age of 65.

10. Have you ever passed out (had vasovagal syncope) during or after a previous immunization or blood draw? [all vaccines]

Providers should be aware of the potential for syncope (fainting) associated with vaccination, particularly among adolescents. Appropriate measures should be taken to prevent syncope, and to readily respond to the patient who feels faint. Observe all patients for 15 minutes after vaccination for signs and symptoms that precede syncope, such as weakness, dizziness, sweateness, and pallor. For patients prone to syncope, make sure they are either seated or lying down at the time of vaccination. (If the patient is seated during vaccination, the immunizer should be seated as well, to minimize the risk of SIRVA). If a patient becomes pre-syncope, have them lie flat or sit with head between knees for several minutes; loosen any tight clothing and maintain an open airway; apply cool, damp cloths to the patient's face and neck. Observe the patient until symptoms completely resolve.

11. Have you received any vaccinations in the past 4 weeks? [LAIV, MMR, VAR, YF vaccine, MVA-BN, COVID-19 vaccines]

Patients who were given either LAIV or an injectable live virus vaccine should wait 28 days before receiving another live vaccine. Inactivated vaccines may be given at the same time or at any spacing interval. There is no minimum interval between receiving any COVID-19 vaccine and MVA-BN (Jynneos), regardless of which vaccine is administered first. However, people, particularly adolescent and young adult males, who are recommended to receive both vaccines might consider waiting 4 weeks between COVID-19 and MVA-BN (Jynneos) vaccines.

12. Are you pregnant, or is there a chance you could become pregnant during the next month? [Adenovirus, AVA, CHIKV vaccine, CVD 103-HgR, HPV, MMR, MVA-BN, LAIV, VAR, Ty21a, YF vaccine]

If giving a vaccine under a standing order during pregnancy (i.e., IIV, Tdap, or RSVpreF), refer to standing order. Live-replicating vaccines (Adenovirus, CHIKV vaccine, CVD 103-HgR, LAIV, MMR, VAR, Ty21a, YF vaccine) and HPV are generally contraindicated during pregnancy.^{6,9} Pregnant women are exempt from AVA and MVA-BN until completion of pregnancy. Other vaccines may be considered during pregnancy with a specific provider order. Women who could become pregnant and plan to receive a live-replicating virus vaccine (e.g., MMR, VAR, YF vaccine) should be instructed to avoid pregnancy for 1 month after vaccination.

Vaccine Abbreviations:

- AVA: Anthrax vaccine (Biothrax)
- CHIKV vaccine: Chikungunya vaccine (Ixchiq)
- CVD 103-HgR: Cholera vaccine (Vaxchora)
- Hib: Haemophilus influenzae type b
- HPV: human papillomavirus
- IIV: inactivated influenza
- IPV: inactivated poliovirus
- LAIV: live attenuated influenza
- MenACWY: meningococcal conjugate, quadrivalent, serogroups A, C, W, Y
- MenB: meningococcal serogroup B
- MMR: measles, mumps, rubella
- MVA-BN: Smallpox/mpox vaccine (Jynneos)
- PPSV23: pneumococcal polysaccharide (23-valent)
- RSVpreF: RSV vaccine (Abrysvo)
- RIV: recombinant influenza
- SIRVA: shoulder injury related to vaccine administration
- Td: tetanus/diphtheria toxoids
- Tdap: tetanus toxoid, reduced diphtheria toxoid, acellular pertussis
- Ty21a: oral typhoid
- VAR: varicella
- YF vaccine: yellow fever

1. ACIP General Best Practice Guidelines for Immunization: www.cdc.gov/vaccines/hcp/acip-recs/general-recs/downloads/general-recs.pdf.
 2. Latex in Vaccine Packaging: www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/latex.pdf.
 3. Table of Vaccine Components: www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/exipient-table-2.pdf.
 4. Prevention and control of seasonal influenza with vaccines: Recommendations of the Advisory Committee on Immunization Practices. www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/flu.html.
 5. Measles, mumps, and rubella-vaccine use and strategies for elimination of measles, rubella, and congenital rubella syndrome and control of mumps. MMWR 1998, 47(RR-8).
 6. Prevention of varicella: Recommendations of the Advisory Committee on Immunization Practices. MMWR 2007, 56(RR-4).
 7. Rubin L.G., Levin M.J., Ljungman P. (2014) IDSA Clinical practice guideline for vaccination of the immunocompromised host. *Clinical Infectious Diseases*, 58(3), 309–318
 8. Tomblyn M, Einsele H, et al. 2009. Guidelines for preventing infectious complications among hematopoietic stem cell transplant recipients: a global perspective. *Biology of Blood and Marrow Transplant* 15:1143-1238.
 9. CDC. Guidelines for Vaccinating Pregnant Women, <https://www.cdc.gov/vaccines/pregnancy/hcp-toolkit/guidelines.html>