

TABLE 2. Prevalence During Second Quarter 2023 of Dispensed Weight Loss Agents Among Active Component, U.S. Armed Forces

	No.	Ratio ^a
Total	1,337	104.4
Sex		
Male	575	54.5
Female	762	339.2
Age group, years		
<20	0	0.0
20-24	89	22.3
25-29	190	62.6
30-34	251	117.9
35-39	352	217.1
40-44	269	319.1
45-49	113	358.6
50+	73	470.6
Service		
Army	419	91.3
Navy	515	157.1
Air Force	330	101.7
Marine Corps	73	43.2
Race and ethnicity		
Non-Hispanic White	655	95.9
Non-Hispanic Black	307	147.3
Hispanic	223	93.0
Other	113	86.4
Unknown	39	211.8
Rank		
Junior enlisted (E1-E4)	159	30.2
Senior enlisted (E5-E9)	756	145.7
Warrant officer (WO1-WO5)	27	144.2
Junior officer (O1-O3)	133	101.8
Senior officer (O4-O10)	260	307.0
Unknown	2	247.2
Military occupation		
Combat-related ^b	55	31.2
Armor / motor transport	20	53.2
Pilot / air crew	10	21.8
Repair / engineering	228	62.3
Communications / intelligence	334	120.1
Health care	475	445.4
Other	215	79.7
History of type 2 diabetes		
No history	1,278	100.0
Previous diagnosis	59	2,124.6

^aPrevalence ratio per 100,000 individuals.

^bInfantry/artillery/combat-engineering/armor.