

TABLE 4b. Demographic and Clinical Characteristic Associations with Long-Term Follow-up Presence of Chronic Pain Diagnosis

Chronic Pain Diagnosis	No		Yes		<i>p</i> -value
	No.	%	No.	%	
Total, <i>n</i>	103 ^a		40 ^a		
Recruit status					0.236
No	89	86.4	38	95.0	
Yes	14	13.6	2	5.0	
Modified Rankin Scale nadir, acute phase					0.080
>3	39	37.9	23	59.0	
0–1	27	26.2	6	15.4	
2	37	35.9	10	25.6	
Pain present, acute phase					<0.001
No	37	35.9	1	2.5	
Yes	66	64.1	39	97.5	
Neuropathic pain treatment					<0.001
No	73	70.6	7	17.5	
Yes	30	29.4	33	82.5	
Results of electrodiagnostic testing ^b					0.035
Total, <i>n</i>	50 ^b		23 ^b		
Axonal, mixed	9	18.0	10	43.5	
Demyelinating	27	54.0	11	47.8	
Normal	14	28.0	2	8.7	

Abbreviations: No., number; *n*, number; CIDP, chronic inflammatory demyelinating polyneuropathy; FND, functional neurologic disorder.

^aLong-term follow-up cases, excluding those with other diagnosis: CIDP, FND, other neurological diagnosis.

^bLong-term follow up cases with electrodiagnostic results, excluding cases with other diagnosis: CIDP, FND, other neurological diagnosis.