

TABLE. Influenza Percent Positivity by Geographic Region of GEIS-funded Laboratory Specimens and WHO GISRS, September 28, 2025–February 7, 2026

Geographic Region	GEIS ^a	GISRS	<i>p</i> -value ^b
North America	13.6	15.5	<0.01
A/not-subtyped	0.0	11.7	
A/H1N1	0.9	0.4	
A/H3N2	10.4	2.5	
B	1.1	0.9	
South America	19.4	10.4	0.33
A/not-subtyped	0.4	3.6	
A/H1N1	1.7	1.7	
A/H3N2	11.9	3.6	
B	1.6	1.5	
West Africa	16.1	19.2	<0.01
A/not-subtyped	0.2	3.6	
A/H1N1	1.8	3.8	
A/H3N2	13.2	11.0	
B	0.8	0.8	
East Africa	17.0	13.1	0.95
A/not-subtyped	7.3	0.8	
A/H1N1	1.7	4.1	
A/H3N2	2.9	7.0	
B	5.1	1.3	
Europe	18.2	24.3	<0.01
A/not-subtyped	2.9	18.5	
A/H1N1	1.7	1.3	
A/H3N2	13.3	4.4	
B	0.0	0.2	
Middle East	29.3	20.5	<0.05
A/not-subtyped	0.0	4.0	
A/H1N1	1.3	2.0	
A/H3N2	17.3	13.3	
B	0.5	1.2	
Asia	30.4	21.2	0.99
A/not-subtyped	10.4	0.1	
A/H1N1	1.1	0.3	
A/H3N2	11.3	20.0	
B	7.3	0.7	

Abbreviations: GEIS, Global Emerging Infections Surveillance; WHO, World Health Organization; GISRS, Global Influenza Surveillance and Response System; DOW, Department of War.

^aInfluenza A, B co-infections included in calculation of overall percent positivity by region for DOW GEIS surveillance.

^b*p*-value corresponding to trend comparisons using dynamic time warping analysis